

PATIENT INFORMATION AND MEDICAL HISTORY

Name: _____ SSN: _____ - _____ - _____

Full Address: _____ City _____ State _____ Zip _____

Cell Phone: _____ Alternate Phone: _____

Date of Birth: //19 Age: _____ Email: _____

HISTORY – Please circle if you have or have had :

- Diabetes
- Sensitive to anesthetic/lidocaine
- Blood clots/thrombus formation
- Lyme disease
- Heart problems or family history of heart problems?
- Hypertension
- Photosensitive disorder
- Autoimmune illness of any type (example: lupus or other)
- History of Bell's Palsy
- Blood or lymphatic disorder or cancer?**

Are you under the care of a physician? If yes, name: _____

Current/Recent medications: _____

Medical illness _____

IF YES, EXPLAIN

- Keloid scars Yes No
- Hives Yes No
- Hypersensitivity to skin products Yes No
- Skin infections Yes No
- Photo sensitizing substances Yes No
- Laser work of any type Yes No

Complications from previous sclerotherapy treatments?

Are you pregnant or breastfeeding? Yes / No

Allergies of any kind _____

I ATTEST THE ABOVE INFORMATION TO BE TRUE, KNOWING MY PROVIDER RELIES ON THIS INFORMATION TO PROVIDE SAFE AND EFFECTIVE TREATMENT. FOR THE SERVICES RECEIVED TODAY, I AGREE TO PAY \$ _____.

Per credit card merchant agreement, I agree to pay the above amount to my credit card. I understand that payment for services rendered is due at time of service. In the event of default, I promise to pay legal interest (18% annually) on the indebtedness, together with such collection costs and reasonable attorney fees as may be required to effect collection of this note.

Adult Signature _____ Date _____

Sclerotherapy Consent Form

This form is designed to provide you with the information you need to make an informed decision regarding sclerotherapy treatment. If you have any questions or do not understand the potential risks, please do not hesitate to ask.

What is sclerotherapy?

Sclerotherapy is a popular method of eliminating varicose veins and superficial telangiectasias (“spider veins”). A solution, called a sclerosing agent, is injected into the veins to break down the vein wall.

Does sclerotherapy work for everyone?

The majority of individuals who receive sclerotherapy treatment will be cleared of their varicosities or have visible improvement. However, there is no guarantee that sclerotherapy will be effective in every case. Approximately 10% of patients who undergo sclerotherapy have poor to fair results. (“Poor results” indicating that the veins have not disappeared after six treatments.) In extremely rare cases, the patient’s condition may become worse after sclerotherapy treatment.

How many treatments will I need?

The number of treatments needed to clear or improve varicose veins differs for each person. The number of treatments can range from one to six, with the average number being three or four. Individual veins usually require one to three treatments.

What are the most common side effects?

The most common side effects experienced with sclerotherapy are:

- 1) Itching: You may experience mild itching along the vein route. This normally lasts 1-2 days.
- 2) Transient Hyperpigmentation: Approximately 30% of all patients who undergo sclerotherapy notice a light brown discoloration after treatment. Nearly all patients notice a darkening of the vein immediately after the procedure. In rare instances the darkening may persist for 4 to 12 months.
- 3) Sloughing: Sloughing occurs in less than 3% of patients who have received sclerotherapy. Sloughing consists of a small, slowly healing ulceration at the injections site. A blister may form, open, and become ulcerated. The scar that follows should return to a normal color.
- 4) Allergic Reactions: Very rarely a patient may have an allergic reaction to the sclerosing agent. This risk is greater in patients who have a history of allergic reactions.
- 5) Pain: A few patients may experience moderate to severe pain and some bruising around the injection site. The veins may be tender to touch after the treatment and an uncomfortable sensation may be felt along the vein route. This pain is usually temporary lasting 1 to 7 days.

What are the other side effects?

Other side effects include a burning sensation during injection of the solution, neovascularization (the temporary development of new, tiny blood vessels), transient phlebitic reactions (temporary swelling of the vein may cause the ankle to swell), temporary superficial blebs (similar to hives), and very rarely wound infection, poor healing, or scarring.

Phlebitis is a very rare complication, seen in approximately 1 out of every 1,000 patients who have been treated for veins greater than 3 or 4 mm in diameter. The dangers of phlebitis include the possibility of pulmonary embolus (a blood clot in the lungs) and postphlebitic syndrome, in which the blood is not carried out of the legs, resulting in permanent swelling of the legs.

What are the possible complications if I do not have sclerotherapy?

In cases of large varicose veins (greater than 3 to 4 mm in diameter), spontaneous phlebitis and/or thrombosis may occur with the associated risk of possible pulmonary emboli. Additionally, large skin ulcerations may develop in the ankle region of patients with long-standing varicose veins with underlying venous insufficiency. Rarely these ulcers may hemorrhage or become cancerous.

Are there other types of procedures to treat large varicose veins and telangiectasias?

Yes, there are other procedures, such as vein stripping and ligation that may be used to treat varicose veins. This generally requires a 1 to 3 week hospital stay and is performed under general anesthesia. Risks of vein stripping or ligation include permanent nerve paralysis in up to 30% of patients and possible pulmonary emboli, infection, and permanent scarring. General anesthesia has some associated risks, including the possibility of serious harm, paralysis, brain damage, and death.

If you notice any type of adverse reaction, please call us immediately.

By initialing I acknowledge that I have received a copy of this sclerotherapy information and consent form.

By signing below, I acknowledge that I have read and the sclerotherapy information and consent form and that the doctor has adequately informed me of the risks of sclerotherapy treatment, alternative methods of treatment, and the risks of not treating my condition. I hereby consent to sclerotherapy treatment to be performed by Dr. Anouche Moshari Roberts, MD.

Patient Signature

Date

Witness

Post Sclerotherapy Instructions

- Wear compression hose overnight on the day of treatment, then only when up and around for the next 5 days
- The veins will look worse before they look better. Bruising is normal.
- Occasionally, there is a hard knot that may be tender. This is normal and will resolve. You can take extra strength Tylenol to relive discomfort.
- Occasionally, there is a brownish discoloration after injections. This is temporary and will fade over time. It can take several months to fade. Avoid sun exposure in these areas until they fade.
- Follow up in 4-6 weeks for next treatment.
- You can resume exercise activities in after 48 hours, light exercise is okay.

If you have any questions, please call Dr Roberts at 703-927-0044